REVIEW OF THE PELAGIC WET-FISHERIES FOR THE 1967–68 SEASON

Anchovy

The 1967–68 season produced one of the smallest wet-fish catches on record. A moratorium on the taking of sardines forced the already low sardine landings even lower. The anchovy fishery, expected to take up the slack in landings as it has in the past two seasons, failed to develop in southern California, and landings in central California remained below the previous year. Pacific mackerel landings were the lowest since Department of Fish and Game records began in 1925. Jack mackerel landings were the lowest in ten years.

Landings for the calendar year, 1967, reveal a slight increase over the previous two years, but the anchovy catch for reduction, made late in the 1966–67 season, accounts for the increase. If the present rate of landing continues, 1968 will be a record low year for wet-fish landings (Table 1).

Vessels continued to leave the wet-fish fleet, and by June of 1968 it consisted of 21 large purse seiners (60 ft. and over), 12 small purse seiners, and 20 lampara boats, a loss of 4 large and 1 small purse seiners since the 1966–67 season. One large and two small purse-seiners fished in central California, and 15 lampara boats fished Monterey Bay. The remainder of the fleet fished southern California waters.

Sardine

A moratorium on the take of sardines in California, effective June 7, 1967, limited the catch to mixed loads with no more than 15% sardines. A total of 41 tons were sold to the fresh-fish markets and brought up to \$500 per ton primarily for use as dead bait. Canners processed 30 tons.

In Baja California the sardine catch was up from last season, at 27,657 tons with most of the catch made in southern Baja California waters (Table 2).

TABLE 1 LANDINGS OF PELAGIC WET-FISHES IN CALIFORNIA IN TONS, 1967–1968

Year	Sar- dines	Ancho- vies	Pacific Mack- erel	Jack Mack- erel	Herring	Squid	Total
1963	3,566	2,285	20,121	47,721	315	5,780	79,788
1964	6,569	2,488	13,414	44,846	175	8,217	75,709
1965	969	2,866	3,525	33,333	258	9,310	50,288
1966	443	31,140	2,314	20,431	121	9,512	63,961
1967	74	34,805	583	19,090	136	9,801	64,489
			1		1		

The California Fish and Game Commission again authorized a 75,000 ton fishery for reduction beginning September 15, 1967, and ending May 15, 1968. The fishery failed to develop in southern California, with only 852 tons landed in the southern area (south of Pt. Conception). Low world prices for fish meal and oil and the resulting low prices paid to fishermen for their catch (\$14-15 per ton), plus a lack of large fish schools close to port combined to inhibit the fishery.

Fishermen in the northern area (north of Pt. Conception) fared much better, delivering 5,651 tons to reduction plants on Monterey Bay, mostly in October, November, and December. This total, however, failed to equal the 8,021 ton catch of the previous season (Table 3).

Reported live-bait catches were below previous 2–3 years (Table 4). The lack of reports from two large operators, who reported in past years, make this total minimal compared to past reports.¹ As in the past, Los Angeles-Long Beach Harbor accounted for about 80% of the take.

Mackerel

Jack and Pacific mackerel catches continued to decline. Pacific mackerel landings of 691 tons were the lowest in the recorded history of the fishery. Jack mackerel landings, although higher than ten years ago, were the lowest since that time. Most of the 18,712 tons taken were caught at Tanner and Cortes Banks. as were some of the Pacific mackerel (Table 5.)—Robson A. Collins, California Department of Fish and Game.

¹ Live-bait catch records are derived from voluntary reports made to the Department of Fish and Game by each fisherman. Not all fishermen report, so these figures should be regarded as minimal.

TABLE 2 SARDINE CATCH IN TONS, 1963–64 THROUGH 1967–68 (Period June Through The Following May)

Season	California	Baja California	Total
1963-64	2,942	18,384	21,362
1964-65	6,103	27,120	33,223
1965-66	962	22,247	23,109
1966-67	344	19,529	19,873
1967-68*	71	27,036	28,007

* Preliminary.

TABLE 3

ANCHOVY LANDINGS FOR REDUCTION IN THE SOUTHERN AND NORTHERN PERMIT AREAS 1965-66 Through 1967-68

Season	Southern Permit Area	Northern Permit Area	Total
965-66*	16,468	375	16,843
1966-67†	29,589	8,021	37,610
.967-68‡	852	5,651	6,503
fotal	46,909	14,047	60,956

Seasons

* October 15, 1965 through April 30, 1966. † October 1, 1966 through April 30, 1967. ‡ September 15, 1967 through May 15, 1968.

TABLE 4

COMMERCIAL LANDINGS AND LIVE BAIT CATCH OF ANCHOVIES IN TONS 1963 THROUGH 1967

Year	Reduction	Other Commercial	Live Bait	Total
1963	0	2,285	$\begin{array}{c} 4,442\\ 5,191\\ 6,148\\ 6,691\\ 5,387\end{array}$	6,727
1964	0	2,488		7,679
1965	170	2,866		9,184
1966	27,335	3,705		37,731
1967	32,349	2,455		40,191

TABLE 5

JACK AND PACIFIC MACKEREL CATCH IN TONS, 1963--64 THROUGH 1965-66

(Period May Through April)

Season	Jack Mackerel	Pacific Mackere
1963-64	42,038	17,105
1964-65	39,557	12,437
1965-66	33,831	3,794
1966-67	22,889	2,037
1967-68	18,712	691

PUBLICATIONS

1 July 1966-30 JUNE 1968

- Aasen, Kenneth D. 1967a. Review of the pelagic wet fisheries for the 1963-64, 1964-65, 1965-66 seasons. Calif. Coop. Ocean. Fish. Invest., Rept., 11:21-22.
- -1967b. Summary of the 1963 and 1964 southern California inshore bait fishery. Calif. Fish and Game, 53(1): 28-34. Ahlstrom, Elbert H. 1967. Co-occurrences of sardine and an-
- chovy larvae in the California Current region off California and Baja California. Calif. Coop. Ocean. Fish. Invest., Rept., 11:117-135.
- Ahlstrom, Elbert H., J. L. Baxter, J. D. Isaacs and P. M. Roedel. 1967. Report of the CalCOFI Committee. Calif. Coop. Ocean. Fish. Invest., Rept., 11:5-9.
- Allen, George H., and Patrick O'Brien. 1967. Preliminary experiments on the acclimatization of juvenile king salmon, Oncorhynchus tshawytscha to saline water mixed with sewage pond effluent. Calif. Fish and Game, 53(3): 180-184.
- Alvariño, A. 1965a. Distributional atlas of Chaetognatha in California Current region. CalCOFI monthly cruises of 1954 and 1958. CalCOFI Atlas, (3): 1-291.
- -1965b. Zoogeography of the Sea of Cortes: Chaetognatha, Siphonophorae and Medusae. Second Mexican Cong. Oceanogr. (In Spanish) (Abstr.)
- -1966. Zoogeography of California: Chaetognatha. Soc. Mexicana Hist. Nat., Rev., 27(2): 236-240; also, Univ. Calif. SIO Contrib., 1967 (2139).

- -1967a. A new Siphonophorae, Vogtia kuruae, n. sp. Pac. Sci., 21(2): 236-240; also, Univ. Calif. SIO Contrib., 1967 (2109)
- -1967b. The Chaetognatha of the NAGA Expedition (1959-1961) in the South China Sea and the Gulf of Thailand. Pt. 1. Systematics. Univ. Calif. SIO NAGA Rept., 4(2): 1 - 197
- 1967c. Evolution in Chaetognatha. Distribution and morphology. Third Mexican Cong. Oceanogr. (In Spanish) (Abstr.)
- -1967d. Bathymetric distribution of Chaetognatha, Siphonophorae, Medusae and Ctenophorae off San Diego, California. Pac. Sci., 21(4): 474-485; also, Univ. Calif. SIO Contrib., 1967 (2171).
- -1967e. Chaetognatha, Siphonophorae and Medusae in the Equatorial Atlantic Ocean, off the Amazon Estuary. Int. Symp. Coastal Lagoons, Mexico City, Nov., 1967. (Abstr.)
- -1968. Two new Calycophorae, Siphonophorae. Pac. Sci.,
- 22(3): 340-346. Baxter, John L. 1966. Anchovy population size. Calif. Dept.
- Fish and Game, MRO Ref., (66-21): 1-2. -1967. Summary of biological information on the northern
- anchovy Engraulis mordax Girard. Calif. Coop. Ocean. Fish. Invest., Rept., 11:110-116.
- -(Editor). 1967. Symposium on anchovies, genus Engraulis. Calif. Coop. Ocean. Fish. Invest., Rept., 11:27-139.
- Beeman, Robert D., and John A. McGowan. 1968. Ophistobranch mollusks from California, Pt. 2. The Order Anaspidea; Thecosomata and Gymnosomata. Veliger, 3, Suppl.: 86-129.
- Benson, A. A. 1967a. On the orientation of lipids on chloroplasts and cell membranes. (Transl. from Japanese by I. Shibuya) Kagaku to Seibutsu (Chem. and Biol.), 5:85-92.
- -1967b. Carbon reduction cycle (in photosynthesis), p. 187-189. In R. J. Williams and E. M. Lansford, Jr. (eds.) Encyclopedia of biochemistry. Reinhold Pub. Co., New York.
- -1967c. Lipid function in plant membranes. Collog. 10, Seventh Int. Cong. Biochem., Tokyo, 3: 525-526. (Abstr.)
- Benson, A. A., and M. R. Atkinson. 1967. Choline sulfate and phosphate in salt excreting plants. Fed. Proc., 26(1): 394. (Abstr. 772)
- Biale, J. B., S. F. Yang and A. A. Benson. 1966. Lipids in plant mitochondria. Fed. Proc., 25(2): 405. (Abstr. 1214) Brinton, Edward. 1966. Remarks and euphausiacean phylogeny.
- Mar. Biol. Ass., India, Symp. on Crustacea, Pt. 1: 255-259. -1967a. Distributional atlas of Euphausiacea (Crustacea)
- in the California Current region, Part 1. CalCOFI Atlas, (5): 1-275.
- -1967b. Vertical migration and avoidance capability of Euphausiids in the California Current. Limnol. Oceanogr., 12(3):451-483.
- -1967c. On the naturalness of change in marine populations. Ass. Fish. Technol. India, Bull.
- Brinton, Edward, and K. Gopalakrishnan. 1968. Preliminary study of the distribution of Euphausiacea in the Indian Ocean. Nat. Inst. Sci. India (Biol.), Proc.
- Brown, Daniel M. 1967. Surface water temperatures at shore stations. U.S. West Coast, 1966. Univ. Calif. SIO Ref., (67 - 9)
- Carlucci, A. F., and S. B. Silbernagel. 1966a. Bioassay of sea water. 2. Methods for the determination of concentrations of dissolved vitamin B₁ in sea water. Can. J. Micorbiol, 12: 1079 - 1089.
- -1966b. Bioassay of sea water. 3. Distribution of vitamin B12 in the northeast Pacific Ocean. Limnol. Oceanogr., 11: 642-646
- -1967a. Bioassay of sea water. 4. The determination of dissolved biotin in sea water using ¹⁴C uptake by cells of Amphidinium carteri. Can. J. Microbiol., 13: 979-986.
- -1967b. Determination of vitamins in seawater. I.B.P. Symp., Chemical environment in the aquatic habitat, N. V. Noord-Hollandsche Vitgevers Maatshappij, Amsterdam, Proc., 239 - 244.
- Carlucci, A. F., and J. D. H. Strickland. 1968. The isolation, purification and some kinetic studies of marine nitrifying bacteria. J. Exper. Mar. Biol. Ecol., 2(2): 156-166.
- Cox, Charles S. 1966. Energy in semidiurnal internal waves. Symp. Mathematical-Hydrodynamical Invest. of Physical